

Statin use in Persons with Diabetes (SUPD)

DENOMINATOR: Patients 40-75 years old who were dispensed at least 2 diabetes medication fills during the measurement year

NUMERATOR: Patient dispensed at least 1 statin medication of any intensity during the measurement year using Medicare Part D benefit.

Statin Medications: Generic and Brand Names			
Atrovastatin (Lipitor, Caduet)	Lovastatin (Altoprev)	Pravastatin (Pravachol)	Simvastatin (Zocor, Vytorin)
Fluvastatin (Lescol XL)	Pitavastatin (Livalo, Zypitamag)	Rosuvastatin (Crestor, Ezallor, Roszet)	

Tips for Coding

- To exclude patients who cannot tolerate statin medications, a claim must be submitted annually using the appropriate ICD-10-CM code
- Providers may use virtual care to confirm and document the exclusion diagnosis in the medical record. They should then bill the non-reimbursable HCPCS code G9781 for \$0.01 with the applicable ICD-10 code attached to process the claim and remove the patient from the Star measure.
- These codes are intended to close Star measure gaps and do not apply payment or reimbursement. Only the codes below will **EXCLUDE** the patient from the SUPD measure:

Exclusion Condition	ICD-10-CM Code
Cirrhosis	K70.30, K70.31, K71.7, K74.3, K74.5, K74.60, K74.69
End Stage Renal Disease (ESRD)	I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.5, Z99.2
Pregnancy and/or Lactation	Numerous > 1k
Polycystic Ovarian Syndrome (PCOS)	E28.2
Pre-diabetes	R73.03
Other abnormal blood glucose	R73.09
Myopathy, drug induced*	G72.0
Myopathy, other specified*	G72.89
Myopathy, unspecified*	G72.9
Myositis, other*	M60.80, M60.819, M60.829, M60.839, M60.849, M60.859, M60.869, M60.879
Rhabdomyolysis*	M62.82
Hospice	Z51.5, along with terminal diagnosis

**The condition the code refers to does not necessarily need to occur in the same year the code was billed. The member's medical chart should reflect "history of"*

Tips for success

- Educate patients on the importance of statin medications for patients with diabetes age 40-75, regardless of LCL levels
- Discuss with patients the importance of taking their medication as prescribed
- Instruct patients to fill prescriptions using their Medicare Part D pharmacy benefit
- Remind patients to contact their practitioner if experiencing medication adverse effects